
Decline of the African lungfish (*Protopterus aethiopicus*) in Lake Victoria (East Africa)

Kees (P. C.) Goudswaard¹, Frans Witte¹ and Lauren J. Chapman²

¹Institute of Evolutionary and Ecological Sciences, University of Leiden, The Netherlands; ²Department of Zoology, University of Florida, Gainesville, Florida, USA, and Wildlife Conservation Society, 185th Street and Southern Blvd., Bronx, New York, USA

Abstract

Catch and effort data for the period 1973–1990 demonstrate a dramatic decline of lungfish in the Tanzanian waters of Lake Victoria. Bottom trawl catches in the Mwanza Gulf showed a decline in catch rates from 67.5 kg h⁻¹ in 1973 to 5.5 kg h⁻¹ in 1986. Trawling of commercial vessels in the Speke Gulf revealed a decline in lungfish catches from 1.3 kg h⁻¹ in 1986 to 0.07 kg h⁻¹ in 1990. The development of anoxia in the deeper waters of Lake Victoria, the algal blooms, and the decline of water transparency, all associated with eutrophication, are not likely to have contributed to the decreased catch rate. However, the lungfish decline may reflect the interaction of overexploitation by the fishery and a low level of Nile perch predation that restricts lungfish to wetland refugia. We suggest that this may have been reinforced over the past few decades by large-scale conversion of wetlands to agricultural land and harvesting of nest-guarding male lungfish leading to decreased recruitment of young.

Key words: habitat deterioration, predation, Nile perch, over exploitation

Résumé

Les données sur les prises et les efforts de capture pour la période qui va de 1973 à 1990 montrent un déclin spectaculaire des protoptères dans les eaux tanzaniennes du lac Victoria. Les prises faites au chalut dans le golfe de Mwanza montraient une réduction des taux de prises de 67,5 kg/h en 1973 à 5,5 kg/h en 1986. Les bateaux commerciaux qui pêchent au chalut dans le golfe de Speke ont rapporté un déclin de leurs prises de protoptères de

1,3 kg/h en 1986 à 0,07 kg/h en 1990. L'extension de l'anoxie dans les eaux profondes du lac Victoria, la profusion d'algues et la diminution de la transparence de l'eau, qui sont toutes associées à l'eutrophisation, sont peu susceptibles d'avoir contribué au déclin du taux de prises. Le déclin des protoptères peut cependant refléter l'interaction entre la surexploitation par la pêche et un faible taux de prédation de la perche du Nil qui repousse les protoptères vers les refuges marécageux. On suggère que ce phénomène a peut-être été renforcé ces dernières décennies par la transformation à grande échelle de terrains marécageux en terres agricoles et par le fait que les prélèvements de protoptères mâles qui gardent les nids ont entraîné une réduction du nombre de jeunes qui grandissent.

Introduction

The four species of African lungfish of the genus *Protopterus* are well-known as obligatory air breathers and for their ability to survive temporary, and sometimes extended, desiccation of their habitat (Smith, 1931; Johnels & Svensson, 1954; Curry-Lindahl, 1956; Greenwood, 1986). The African lungfish tend to inhabit shallow inshore waters, including the dense interior of swamps and marshes, although larger individuals can be found offshore in some lakes, such as Lake Victoria (Johnels & Svensson, 1954; Curry-Lindahl, 1956; Greenwood, 1966; Greenwood, 1986; Chapman *et al.*, 1996). Their snake-like appearance and the strong taste of their flesh, places them in a particular category for fishermen and consumers. In the Lake Victoria basin, the African lungfish, *P. aethiopicus* Heckel (1851), is locally either highly appreciated or strongly disliked. For example, for the Luo, who live south of the Nyanza Gulf (Fig. 1), the lungfish was particularly popular as a food fish (Graham, 1929;

Correspondence: Frans Witte, Institute of Evolutionary and Ecological Sciences, University of Leiden, Postbox 9516, 2300 RA Leiden, The Netherlands. E-mail: Witte@rulsfb.leidenuniv.nl




Fig 1 Study sites in Lake Victoria. Asterisks indicate the five sampling stations of M.V. Kiboko. Hatched areas in the Mwanza Gulf indicate fishing areas of M.V. Mdiria. Hatched areas in the Speke Gulf indicate a commercially trawled area

P. C. Goudswaard, pers. obs.), and in Lake Nabugabo, Uganda, the lungfish contributed most to the biomass of fish landed in the 1960s (Cambridge Nabugabo Biological Survey, unpubl. obs.). In contrast, the Sukuma, who live around the Mwanza and Speke Gulfs disliked this fish.

Fishing for lungfish in Lake Victoria has been practised for a long time. However, a limited number of specialised fishermen targeted this species. Until the 1950s, the primary species exploited by most fishermen was the tilapiine cichlid *Oreochromis esculentus* (Graham, 1929). Gill nets and long lines for lungfish were placed along the lakeshore, close to the vegetation zone, mostly papyrus stands, where the best catches were made. In the long line fishery, hooks were baited

with the flesh of freshwater mussels or haplochromines.

Lungfish were usually sold as sun-dried products. Both in 1928 (Graham, 1929) and in the early 1980s (P. C. Goudswaard, pers. obs.) catches comprised only a few fish per fisherman per night. Storage of dried lungfish enabled the fishermen to collect sufficient fish to make a deal with a trader who transported the dried product to market. From the southern coast of the lake between Speke and Emin Pasha Gulf, almost all of the dried lungfish was transported to Kenya and the area north of Musoma town where the Luo people live (P. C. Goudswaard, pers. obs.). This type of fishery, without central landing and trading places, makes the collection of catch statistics for lungfish difficult. However, a

lake-wide decline in the lungfish landings between 1970 and 1990 was detected (Ogutu-Ohwayo, 1990a; Ochumba, 1995).

The upsurge in the 1980s of the introduced piscivorous Nile perch, *Lates niloticus* (Linnaeus, 1758) coincided with dramatic changes in the species composition of Lake Victoria (Ogutu-Ohwayo, 1990a; Witte *et al.*, 1992b; Goudswaard & Witte, 1997). Fisheries over-exploitation (Cadwalladr, 1965; Fryer & Iles, 1972; Witte & Goudswaard, 1985) and eutrophication (Seehausen, van Alphen & Witte, 1997; Wanink *et al.*, 2001) also contributed to these changes. In this study, we analyse bottom trawl catches from the period 1973–1990 in southern Lake Victoria to quantify the degree of the decline in lungfish. We discuss the possible impacts of Nile perch predation, over-exploitation and environmental changes on the lungfish stock.

Material and methods

Sources of information can be divided into four categories:

1 Data of commercial operating bottom trawlers were used: (i) M.V. Mdiria (120 hp, trawl headrope 25 m, codend mesh 20 mm [mesh sizes throughout the paper refer to stretched meshes]) of the Freshwater Fisheries Training Institute Nyegezi, aimed at haplochromine cichlids in the Mwanza Gulf (Fig. 1) at depths of 4–18 m. Data were available from 1973–1986. (ii) Three trawlers (58 hp, trawl headrope 17 m, codend mesh 70 mm) aimed at Nile perch in the Speke Gulf (Fig. 1) at depths of 10–20 m. Data from these ships were available from 1986–1990 and were cross-checked at intervals by measuring the complete catch. In both cases, trawl shots generally lasted 2 h. *P. aethiopicus* was bycatch.




Fig 2 Length frequency distribution of lungfish: (a) in long line catches in the Tanzanian waters from 1973 to 1975, and (b) in trawl catches of M.V. Kiboko at five stations in the Mwanza Gulf during 1984–90. The percentage of Nile perch, caught during 1984–90, able to swallow lungfish of a particular length based on mouth gape of Nile perch is indicated by the shaded area under the solid line. The dark shaded area under dotted line indicates the percentage of Nile perch able to swallow lungfish of 50% of their own length. Inset: length frequency distributions of lungfish in 1984 before the Nile perch boom

2 Data from M.V. Kiboko (105 hp, bottom trawl headrope 18 m, codend mesh 20 mm) of the Haplochromis Ecology Survey Team (HEST) and the Tanzania Fisheries Research Institute (TAFIRI). This vessel made trawl shots of 30 min. From October 1984 until July 1990, five stations in the Mwanza Gulf (Fig. 1), with soft mud bottoms and depths ranging from 1–2 m in the south to 35 m in the north, were sampled on a monthly basis.

3 Data collected with long lines and gill nets (3.5'' and 4'' mesh) by the East African Freshwater Fisheries Research Organization (EAFFRO) during the period 1973–1975 in the Tanzanian part of Lake Victoria. The length frequency distribution from 1973–1975 (Fig. 2a) is based upon the long line data, because long lines are less size selective than gill nets.

4 Data collected with metal minnow traps (6.3 mm mesh) and gill nets (25.4, 50.5 and 76.2 mm mesh) from 1995–1999 during surveys of twelve small swamp lagoons, deep in the interior of Lwamunda Swamp surrounding Lake Nabugabo (Fig. 1), a satellite lake of Lake Victoria. The lagoons are separated from the main lake by 150–3000 m fringing swamp, and grade into the *Miscanthidium violaceum* dominated wetland. Seines and dip nets were used to qualitatively sample edge areas.

In the present study, the length of lungfish was measured as total length (TL) in centimetres. No correction was made for damaged tail tips, because missing parts were relatively small. Of fish collected by EAFFRO in 1973–1975, anal length (AL in centimetres; distance from




Fig 3 Lungfish catches in bottom trawls of: (a) M.V. Mdiria in the Mwanza Gulf from 1973 to 1986 with 20 mm codend mesh. (b) M.V. Kiboko in the Mwanza Gulf from 1984 to 1990 with 20 mm codend mesh. (c) commercial trawlers in the Speke Gulf from 1986 to 1990 with 70 mm codend mesh. Note the different scales on the Y-axes

tip of the nose to the anal pore) was recorded. This was converted to TL by: $TL = 1.75 AL + 0.08$ ($n = 27$; $R^2 = 0.9745$).

We estimated relative vulnerability of lungfish to Nile perch predation by relating maximum body depth of lungfish to the mouth gape of Nile perch. The mouth gape (MG) of *Lates niloticus* is related to its total length: $MG = 0.112 \times TL + 0.43$ (Ligtvoet & Mkumbo, 1990 – transformed to centimetres). Body depth of lungfish was calculated from maximum girth (G in centimetres) by considering the fish circular in cross-section ($G = 2\pi R$). We used girth–length relationship: $G = 0.565 \times AL - 1.642$ (Witte & de Winter, 1995). To calculate the fraction

of Nile perch able to swallow lungfish of a certain size, we used all Nile perch above 25 cm TL, caught between 1984 and 1990 at the four shallowest stations in the Mwanza Gulf (Fig. 1). For Nile perch greater than 25 cm, fish often constitutes a major part of the diet (Hamblyn, 1966; Ogari & Dadzie, 1988; Ogutu-Ohwayo, 1990b).

Results

Bottom trawl catches by M. V. Mdiria in the Mwanza Gulf showed a decline in lungfish catch rates from 67.5 kg h^{-1} in 1973 to 5.5 kg h^{-1} in 1986 (Fig. 3a). Catches of M. V. Kiboko from 1984 to 1990 in the same area revealed that




Fig 4 (a) Decrease in number of lungfish at different stations, and (b) increase of average lungfish length at all stations combined, in trawl catches of M.V. Kiboko in the Mwanza Gulf from 1984 to 1990. Error bars indicate SD, numbers indicate number of fish

lungfish continued to decline (Fig. 3b). Trawling of commercial vessels in the Speke Gulf revealed a decline in lungfish catches from 1.3 kg h^{-1} in 1986 to 0.07 kg h^{-1} in 1990 (Fig. 3c).

Bottom trawling at the two deepest stations in front of the Mwanza Gulf, 30–35 and 57 m deep, did not produce any lungfish. Most fish were captured in the gulf, at the three stations shallower than 16 m, while at the 20–25 m deep station, considerably less lungfish were caught (Fig. 4a). Large Nile perch showed a complimentary distribution. Between 1984 and 1990, large Nile perch were most abundant in deeper waters (Fig. 5). A strong decline of lungfish catches with increasing depth was also observed during a trawl survey in 1969–1970 (Kudhongania & Cordone, 1974a,b). While the number of lungfish decreased during the years 1984–1990 (Fig. 4a), the average size increased gradually at all sampling stations in the Mwanza Gulf (Fig. 4b; ANOVA $P < 0.000$; Pearson correlation for means $R = 0.9115$, $P = 0.0043$).

The length frequency distribution of all lungfish caught by M. V. Kiboko in the Mwanza Gulf during 1984–1990 ranged from 42 to 186 cm, 90% of the lungfish were between 90 and 130 cm (Fig. 2b). The length




Fig 5 Mean catch rate of Nile perch >25 and >50 cm TL in bottom trawls of M. V. Kiboko (1984–1990) at different stations in the Mwanza Gulf




Fig 6 Length frequency distribution of lungfish caught in Lwamunda Swamp from 1995–1999 with minnow traps and gill nets

frequency distribution of lungfish from the long line fishery in 1973–1975 ranged from 42 to 164 cm with 90% of the fish between 56 and 126 cm (Fig. 2a). In 1974, gill net fishermen caught fish from 45 to 72 cm ($n = 63$) with 88.9 mm (3.5") mesh nets and from 44 to 93 cm ($n = 45$) with 101.6 mm (4") mesh nets.

Lungfish smaller than 40 cm are very rare in open waters of Lake Victoria (Fig. 2). This is supported by data from a satellite of Lake Victoria, Lake Nabugabo. Data from the Lwamunda Swamp surrounding the satellite Lake Nabugabo (Fig. 1) suggest that smaller lungfish are most abundant within the fringing swamp. The length of lungfish from traps and gill nets in swamp lagoons of the Lwamunda Swamp during 1995–1999 ranged from 18 to 60 cm with 90% of the fish being between 20 and 45 cm TL (Fig. 6). Small lungfish (<10 cm) were dip netted and seined from within the dense *Miscanthidium* vegetation, whereas lungfish greater than 20 cm were more common in the edge areas of the lagoons. Similar observations were made by Graham (1929) and Greenwood (1966), for other parts of Lake Victoria.

Plotting cumulative frequency distributions of Nile perch with a mouth gape that could swallow a lungfish of a certain length (based on its maximum girth), revealed that a relatively small fraction of the lungfish were vulnerable to Nile perch predation in the area fished by trawlers during 1984–1990 (Fig. 2b). It seems like a much larger percentage would have been available to Nile perch in the 1973–1975 period (Fig. 2a), prior to the upsurge of

the predator. However, long lines that are normally used in shallower areas caught these fish.

Discussion

Bottom trawl catches in the Mwanza and Speke Gulfs confirm the decline of lungfish indicated by commercial landing records of the three riparian countries of Lake Victoria (Ogotu-Ohwayo, 1990a; Ochumba, 1995). In the Mwanza Gulf, which was monitored most intensively during the longest period, catches dropped an order of magnitude between 1973 and 1990. This decline may reflect the impact of the fisheries, the impact of Nile perch, decline in food abundance, and/or habitat deterioration.

Impact of the fisheries

Lungfish smaller than 40 cm TL were rarely encountered in trawl catches and in any of the catches with other gear in open waters (Okedi, 1971a; Mosille & Mainoya, 1988). Graham (1929) recorded a size range of lungfish from 61 to 126 cm in the lake and noted that small fish were rarely caught outside vegetated zones. Juvenile lungfish (5–30 cm), were found in matted roots of papyrus, and may be limited to these habitats (Graham, 1929; Greenwood, 1966). Our results indicate that the smallest individuals (<10 cm) live primarily within the dense mats of wetland vegetation, fish of 20–40 cm are also most abundant within the swamps, particularly swamp lagoons; whereas fish larger than 40 cm predominantly occur in the main lake.

The size at first maturity during the 1980s in the Mwanza Gulf was 65–75 cm for males and 75–85 cm for females (Mosille & Mainoya, 1988). Greenwood (1958) found first sexual maturity of females between 65 and 70 cm during the 1950s. The percentage of immature lungfish (<70 cm) in the catches was 22% in the 1973–1975 period, and therefore may have contributed to a modest reduction in stock recruitment. However, between 1984 and 1990, immature fish represented only 2% of the catch, and therefore stock recruitment overfishing was negligible during this time period. Except in the Nyanza Gulf (CIFA, 1982), the specialised fishery on adult lungfish was probably not intensive enough to explain the lake-wide decline of the lungfish stocks, although it may have contributed locally to the decline. Bottom trawling for haplochromines in the Mwanza Gulf included considerable quantities of lungfish until the first half of the 1980s (Kudhongania & Cordone, 1974a; Goudswaard,

1988), and certainly could have impacted lungfish stocks. However, the decline in the Mwanza Gulf continued after 1987, when commercial trawling in this area was abandoned. Thus, trawling cannot be responsible for the decline during this period. This conclusion is supported by low catch rates (c. 1.3 kg h⁻¹) in the Speke Gulf in 1986 compared to the catch rates of 24.4 kg h⁻¹ in 1971 in the same area (Kudhongania & Cordone, 1974a), whereas here the commercial trawl fishery only started in 1986.

Impact of the Nile perch

Habitat overlap of haplochromine cichlids, catfish, and the cyprinid *Rastrineobola argentea* (Pellegrin, 1904) with Nile perch, was one of the major factors determining if, and at what speed, population densities declined after the Nile perch boom (Wanink, 1991; Witte *et al.*, 1992b; Goudswaard & Witte, 1997). There is little habitat overlap between juvenile lungfish and Nile perch. Juvenile lungfish, up to 40 cm, occur primarily in swamps and swamp lagoons, which are less frequently used by Nile perch because of reduced oxygen availability and structural complexity (Schofield & Chapman, 1999, 2000). Therefore, Nile perch predation on juvenile lungfish seems negligible. This is supported by stomach content data on Nile perch of a size most likely to consume small lungfish from lakes Victoria, Kyoga and Albert (Ogotu-Ohwayo, 1994) and Lake Nabugabo (Ogotu-Ohwayo, 1993, 1994; Schofield & Chapman, 1999).

Nile perch greater than 25 cm TL feed to a large extent on fish (Hamblyn, 1966; Ogari & Dadzie, 1988; Ogotu-Ohwayo, 1990b). Piscivorous fish are generally gape-limited predators (Hambright, 1991; Brönmark & Miner, 1992). Based on mouth gape data, a Nile perch of 25 cm could theoretically swallow a lungfish of 40 cm, and the largest Nile perch encountered in the Mwanza Gulf (161 cm TL) should be able to swallow a lungfish of 185 cm. However, this is a very conservative estimate because: (i) the cross-section of lungfish is not circular (see Material and methods); its body is deeper than wide, implying that our calculations, based on mouth gape, overestimate the size of the lungfish that can be swallowed. Moreover, piscivores tend to feed on prey that are smaller than the maximum size possible (Gillen, Stein & Carline, 1981; Hambright, 1991; Brönmark & Miner, 1992). (ii) Nile perch of >25 cm are predominantly caught at depths of more than 12 m and the highest numbers of Nile perch of >50 cm are caught at depths of more than

20 m, where lungfish are less abundant. (iii) The estimate, based on mouth gape, would enable Nile perch to swallow lungfish above its own length. This is unlikely, even though their eel-like body may curl up more easily than that of most other fish. The maximum fish prey size ever recorded for Nile perch approaches 50% of its own length (Ogari & Dadzie, 1988). However, most studies indicate a prey body-length to predator body-length ratio of 20–35% (Hamblyn, 1966; Gee, 1969; Okedi, 1971b). Using the upper limit of 50% of prey to predator body-length implies that the largest Nile perch in the Mwanza Gulf could swallow a lungfish of 80 cm. Consequently, most Nile perch are probably too small to feed on the available size of lungfish in the main lake (Fig. 2b).

Diet data on Nile perch support the idea that they do not feed intensively on large lungfish. Stomach investigations of hundreds of Nile perch in southern Lake Victoria revealed only two adult lungfishes (P. C. Goudswaard, pers. obs.), each eaten by a Nile perch of more than 100 cm. In Lake Kyoga, lungfish is eaten in small quantities by large (>120 cm) Nile perch (Ogututu-Ohwayo, 1990b). In the Nyanza and Mwanza Gulf, lungfish was reported as rare in Nile perch stomachs (Ogari & Dadzie, 1988; Mkumbo & Ligtvoet, 1992).

Comparing the length frequency distribution of 1984 with the data from 1984 to 1990 collected with the same gear in the same areas of the Mwanza Gulf, shows that 40–80 cm lungfish were already rare in 1984 before the Nile perch increased (Fig. 2b). This indicates that the length frequency distribution as observed in the Mwanza Gulf has not been strongly affected by Nile perch predation.

Summarizing, an effect of Nile perch predation on the lungfish stocks cannot be excluded, but evidence suggests that it cannot alone account for the decline in lungfish catches. However, it is possible that Nile perch predation pressure on the few small individuals that venture out of the wetlands and on the occasional large lungfish may reinforce use of wetland refugia. In addition, Nile perch predation in the early 1980s when the upsurge began may have contributed to the shift in size distribution observed by 1984 (compare Fig. 2a with inset in Fig. 2b). Although, as explained above, habitat differences could also account for this.

Impact of food abundance

The diet of adult and sub-adult lungfish consists primarily of molluscs, mainly the gastropods of the genus

Bellamyia and *Melanoides* (Corbet, 1961). These snails were abundant in the shallow waters of Lake Victoria (Okedi, 1990). In the second half of the 1980s, an increase in *Bellamyia* was noticed during bottom trawling in shallow waters of Mwanza Gulf, and these snails then often formed the bulk of the catch (P. C. Goudswaard, pers. obs.). This increase may reflect a decline in the amount of molluscivorous fish in the lake (Witte *et al.*, 1992a). In the same period, mass snail mortalities were observed in the deeper parts of Speke and Mwanza Gulfs, which may have been caused by deoxygenation of deeper water layers (Kaufman, 1992; Hecky, 1993; Wanink *et al.* 2001). Despite these mass kills, *Bellamyia* remained very abundant in the habitat of large lungfish, and it is unlikely that food shortage has played a role in the lungfish decline.

Impact of habitat deterioration

Over the past three decades many dramatic environmental changes have occurred in Lake Victoria. Most striking have been an increase in algal blooms, a decrease in water transparency, and a decrease in dissolved oxygen concentration in deeper waters (Ochumba & Kibaara, 1989; Kaufman, 1992; Witte *et al.*, 1992a; Hecky, 1993; Mugidde, 1993; Wanink *et al.*, 2001). However, due to the dependence of lungfish on atmospheric oxygen, their ability to live in swamps, and their reliance on nonvisual cues for feeding (Greenwood, 1986) the lungfish is one of the few fish species not likely to be affected by these environmental changes.

Deterioration of the lungfish breeding-habitat may be a key environmental factor affecting the lungfish population. The lungfish performs parental care in which the male prepares a nest in a pit or a hole, where he guards the eggs and young for nearly 8 weeks (Greenwood, 1958, 1966; 1986). More than one female may spawn in a nest. The nature of the nest is variable, including nests excavated in matted roots of papyrus, among plants in shallow grass-swamps, and in the sand beneath a fallen tree (Greenwood, 1958, 1966; 1986). However, in general most nests seem to occur in wetlands.

The human population bordering Lake Victoria is among the densest in East Africa and shows an annual growth of 3.8% (Bootsma & Hecky, 1993). To feed this population, large strips of swamp near the lakeshore and along in-flowing rivers have been converted to agriculture (Balirwa, 1998). These cultivated areas usually reach the water where all vegetation is cleared; making

these places unsuitable for lungfish to breed. However, this may not be the only explanation for the decline of lungfish because the species declined dramatically in Lake Nabugabo (Ogutu-Ohwayo, 1993; Chapman *et al.*, 1996), without coincident loss of wetland (L. J. Chapman, pers. obs.).

Between 1982 and 1990, we found two nest guarding lungfish in the Mwanza Gulf area. Both nests were in disorder within a few days after discovery, apparently after being raided. Local farmers and fishermen told on several occasions that, when they discover a lungfish nest, the guarding fish is killed by club or spear and taken for food. Similar observations were made by Wasawo (1959), with respect to the dry season burrows. We tentatively suggest that exploitation of nest guarding fish, together with breeding habitat destruction, may have contributed to the population decline through decreased recruitment of young fish. This may also explain the increase in mean weight (ageing of the stock) of lungfish during the 1980s.

To summarise, catch and effort data (bottom trawl) for the period 1973–1990, demonstrate a dramatic decline of lungfish in the Tanzanian waters of Lake Victoria. In contrast to the decline of many other fish species in Lake Victoria, Nile perch predation may not have been a key factor contributing to the dramatic decline of the lungfish. In addition, the development of anoxia in the deeper waters of Lake Victoria, the algal blooms, and the decline of water transparency, are not likely to have affected the lungfish population in a negative way. However, the decline may reflect the interaction of local overexploitation and a low level of Nile perch predation that restricts lungfish to wetland refugia. We suggest that this may have been reinforced over the past few decades by large-scale conversion of wetlands to agricultural land and harvesting of nest-guarding males leading to decreased recruitment of young lungfish.

Preliminary observations indicate some recovery of lungfish since 1997 (Bugenyi & v.d. Knaap, 1997). The authors attribute this modest resurgence to the extraordinary spread of water hyacinth (*Eichhornia crassipes*), which invaded the lake in 1989 (Njuguna, 1991). The extensive mats of this weed along the lake's shore extend ecotonal refugia, thus potentially increasing availability of lungfish habitat. However, if lungfish really benefit from the water hyacinth infestation of the lake, it is potentially short-lived, because a dramatic decline of water hyacinth has occurred through Lake Victoria since 1999.

Acknowledgements

We are grateful to Mr. E. F. B. Katunzi and the other staff of the Tanzania Fisheries Research Institute for the hospitality and support. We are indebted to Mr. R. Mapunda, former director of the Freshwater Fisheries Training Institute, who kindly provided the catch data of M. V. Mdiria, and to Mr. Maketoshi who gave us the catch records from commercial trawlers in the Speke Gulf. The crew of M. V. Kiboko is thanked for their dedication in collecting data often under difficult circumstances. Drs. C. D. N. Barel and J. H. Wanink are acknowledged for their comments on earlier drafts of the paper. This research was part of the Haplochromis Ecology Survey Team (HEST) and was financially supported by the section for Research and Technology of The Netherlands Minister of Development Cooperation, the Wildlife Conservation Society, USA (L.J.C.), and the National Science Foundation, USA (DEB-9622218 to LCJ).

References

- BALIRWA, J.S. (1998) *Lake Victoria Wetlands and the Ecology of the Nile tilapia, Oreochromis niloticus* Linne. (PhD Thesis) A.A. Balkema, Rotterdam.
- BOOTSMA, H.A. & HECKY, R.E. (1993) Conservation of the Great Lakes: a limnological perspective. *Conserv. Biol.* **7**, 644–655.
- BRÖNMARK, C. & MINER, J.G. (1992) Predator-induced phenotypical change in body morphology in crucian carp. *Science* **258**, 1348–1350.
- BUGENYI, F.W.B. & v.d. KNAAP, M. (1997) Lake Victoria: an example of ACP-EU fisheries research. *EC Fish. Coop. Bull.* **10**, 20–21.
- CADWALLADR, D.A. (1965) The decline in the *Labeo victorinus* Boulenger (Pisces: Cyprinidae) fishery of Lake Victoria and an associated deterioration in some indigenous fishing methods in the Nzoia River, Kenya. *E. Afr. Agric. For. J.* **30**, 249–256.
- CHAPMAN, L.J., CHAPMAN, C.A., OGUTU-OHWAYO, R., CHANDLER, M., KAUFMAN, L. & KEITER, A.E. (1996) Refugia for endangered fishes from an introduced predator in Lake Nabugabo, Uganda. *Cons. Biol.* **10**, 554–561.
- CIFA (1982) *Report of the First Session of the Sub-committee for the Development and Management of the Fisheries of Lake Victoria, 12–14 October 1981, Mwanza, Tanzania*. FAO Fisheries Report 262. FAO, Rome.
- CORBET, P.S. (1961) The food of non-cichlid fishes in the Lake Victoria basin, with remarks on their evolution and adaptation to lacustrine conditions. *Proc. Zool. Soc. Lond.* **136**, 1–101.
- CURRY-LINDAHL, K. (1956) On the ecology, feeding behaviour and territoriality of the African lungfish, *Protopterus aethiopicus* Heckel. *Ark. Zool.* **9**, 479–497.

- FRYER, G. & ILES, T.D. (1972) *The Cichlid Fishes of the Great Lakes of Africa*. Oliver & Boyd, Edinburgh.
- GEE, J.M. (1969) A comparison of certain aspects of the biology of *Lates niloticus* (Linné) in some African lakes. *Rev. Zool. Bot. Afr.* **80**, 244–262.
- GILLEN, A.L., STEIN, R.A. & CARLINE, R.F. (1981) Predation by pellet-reared tiger muskellunge on minnows and bluegills in experimental systems. *Trans. Am. Fish. Soc.* **110**, 197–209.
- GOUDSWAARD, P.C. (1988) A comparison of trawl surveys in 1969/70 and 1984/85 in the Tanzanian Part of Lake Victoria. In: *Report of the Fourth Session of the Sub-committee for the Development and Management of the Fisheries in Lake Victoria, 6–10 April 1987, Kisumu, Kenya (CIFA)*. FAO Fisheries Report 388. FAO, Rome.
- GOUDSWAARD, P.C. & WITTE, F. (1997) The catfish fauna of Lake Victoria after the Nile perch upsurge. *Env. Biol. Fish.* **49**, 21–43.
- GRAHAM, M. (1929) *The Victoria Nyanza and its Fisheries. A Report of the Fishing Survey of Lake Victoria (1927–28)*. Crown Agents for the Colonies, London.
- GREENWOOD, P.H. (1958) Reproduction in the East African lung-fish *Protopterus aethiopicus* Heckel. *Proc. Zool. Soc. London* **130**, 547–567.
- GREENWOOD, P.H. (1966) *The Fishes of Uganda*. The Uganda Society, Kampala.
- GREENWOOD, P.H. (1986) The natural history of African lungfishes. *J. Morph. Suppl.* **1**, 163–179.
- HAMBLYN, E.L. (1966) The food and feeding habits of Nile perch: *Lates niloticus* (Linné) (Pisces: Centropomidae). *Rev. Zool. Bot. Afr.* **74**, 1–28.
- HAMBRIGHT, K.D. (1991) Experimental analysis of prey selection by largemouth bass: role of predator mouth width and prey body depth. *Trans. Am. Fish. Soc.* **120**, 500–508.
- HECKY, R.E. (1993) The eutrophication of Lake Victoria. *Verh. Internat. Verein. Limnol.* **25**, 39–48.
- JOHNELS, A.G. & SVENSSON, S.O. (1954) On the biology of *Protopterus annectens* (Owen). *Ark Zool.* **7**, 131–164.
- KAUFMAN, L. (1992) Catastrophic change in species-rich freshwater ecosystems: the lessons of Lake Victoria. *Bioscience* **42**, 846–858.
- KUDHONGANIA, A.W. & CORDONE, A.J. (1974a) Past trends, present stocks and possible future state of the fisheries of the Tanzania part of the Lake Victoria. *Afr. J. trop. Hydrobiol. Fish.* **3**, 167–181.
- KUDHONGANIA, A.W. & CORDONE, A.J. (1974b) Batho-spatial distribution pattern and biomass estimate of the major demersal fishes in Lake Victoria. *Afr. J. trop. Hydrobiol. Fish.* **3**, 15–31.
- LIGTVOET, W. & MKUMBO, O.C. (1990) Synopsis of ecological and fishery research on Nile perch (*Lates niloticus*) in Lake Victoria, conducted by HEST/TAFIRI. In: *Report of the Fifth Session of the Sub-committee for the Development and Management of the Fisheries in Lake Victoria, 12–14 September 1989, Mwanza, Tanzania (CIFA)*. FAO Fisheries Report 430. FAO, Rome.
- MKUMBO, O. & LIGTVOET, W. (1992) Changes in the diet of Nile perch, *Lates niloticus*, in the Mwanza Gulf, Lake Victoria. *Hydrobiologia* **232**, 79–84.
- MOSILLE, O.I.I.W. & MAINOYA, J.R. (1988) Reproductive biology of the East African Lungfish (*Protopterus aethiopicus*) in Mwanza Gulf, Lake Victoria. *Afr. J. Ecol.* **26**, 149–162.
- MUGIDDE, R. (1993) The increase in phytoplankton primary productivity and biomass in Lake Victoria (Uganda). *Verh. Internat. Verein. Limnol.* **25**, 846–849.
- NJUGUNA, S.G. (1991) Water hyacinth: the worlds worst aquatic weed infests Lakes Naivasha and Victoria. *Swara* **14** (6), 8–10.
- OCHUMBA, P.B.O. (1995) Limnological changes in Lake Victoria since the Nile perch introduction. In: *Impact of Species Changes in African Lakes* (Eds T. J. PITCHER and P. J. B. HART). Chapman & Hall, London.
- OCHUMBA, P.B.O. & KIBAARA, D.I. (1989) Observations on blue-green algal blooms in the open waters of Lake Victoria, Kenya. *Afr. J. Ecol.* **27**, 23–34.
- OGARI, J. & DADZIE, S. (1988) The food of the Nile perch, *Lates niloticus* (L.), after the disappearance of the haplochromine cichlids in the Nyanza Gulf of Lake Victoria (Kenya). *J. Fish Biol.* **32**, 571–577.
- OGUTU-OHWAYO, R. (1990a) The decline of the native fishes of Lakes Victoria and Kyoga (East Africa) and the impact of introduced species, especially the Nile perch, *Lates niloticus* and the Nile tilapia, *Oreochromis niloticus*. *Env. Biol. Fish.* **27**, 81–96.
- OGUTU-OHWAYO, R. (1990b) Changes in the prey ingested and the variations in the Nile perch and other fish stocks of Lake Kyoga and the northern waters of Lake Victoria (Uganda). *J. Fish. Biol.* **37**, 55–63.
- OGUTU-OHWAYO, R. (1993) The effect of predation by Nile perch, *Lates niloticus* L., on the fish of Lake Nabugabo, with suggestions for conservation of endangered endemic cichlids. *Cons. Biol.* **7**, 701–711.
- OGUTU-OHWAYO, R. (1994) *Adjustments in Fish Stocks and in Life History Characteristics of the Nile Perch, Lates niloticus L. in Lakes Victoria, Kyoga and Nabugabo*. PhD Dissertation, University of Manitoba, Winnipeg, Manitoba.
- OKEDI, J. (1971a) *Maturity, Sex Ratio and Fecundity of the Lungfish (Protopterus aethiopicus Heckel) from Lake Victoria*. EAFFRO Annual Report (1970), pp. 17–20.
- OKEDI, J. (1971b) *Further Observations on the Ecology of the Nile perch (Lates niloticus Linné) in Lake Victoria and Lake Kyoga*. EAFFRO Annual Report (1970), pp. 42–55.
- OKEDI, J. (1990) Observations on the benthos of Murchinson Bay, Lake Victoria, East Africa. *Afr. J. Ecol.* **28**, 111–122.
- SCHOFIELD, P.J. & CHAPMAN, L.J. (1999) Interactions between Nile perch, *Lates niloticus*, and other fishes in Lake Nabugabo, Uganda. *Env. Biol. Fish.* **55**, 343–358.
- SCHOFIELD, P.J. & CHAPMAN, L.J. (2000) Hypoxia tolerance of introduced Nile perch: implications for survival of indigenous fishes in the Lake Victoria basin. *Afr. Zool.* **35**, 35–42.
- SEEHAUSEN, O., VAN ALPHEN, J.J.M. & WITTE, F. (1997) Cichlid fish diversity threatened by eutrophication that curbs sexual selection. *Science* **277**, 1801–1811.

- SMITH, H.W. (1931) Observations on the African lungfish, *Protopterus aethiopicus*, and on evolution from water to land environments. *Ecology* **12**, 164–181.
- WANINK, J.H. (1991) Survival in a perturbed environment: the effects of Nile perch introduction on the zooplanktivorous fish community of Lake Victoria. In: *Terrestrial and Aquatic Ecosystems: Perturbation and Recovery* (Ed. O. RAVERA). Ellis Horwood, Chichester.
- WANINK, J.H., KASHINDYE, J.J., GOUDSWAARD, P.C. & WITTE, F. (2001) Dwelling at the oxycline: does increased stratification provide a predation refugium for the Lake Victoria sardine *Rastrineobola argentea*? *Freshwat. Biol.* **46**, 75–85.
- WASAWO, D.P.S. (1959) A dry season burrow of *Protopterus aethiopicus* Heckel. *Rev. Zool. Bot. Afr.* **60**, 65–70.
- WITTE, F. & DE WINTER, W. (1995) Biology of the major fish species of Lake Victoria. In: *Fish Stocks and Fisheries of Lake Victoria: a Handbook for Field Observations* (Eds F. WITTE and W. L. T. VAN DENSEN). Samara Publishing Ltd, Cardigan, Dyfed, UK.
- WITTE, E., GOLDSCHMIDT, T., GOUDSWAARD, P.C., LIGTVOET, W., VAN OIJEN, M.J.P. & WANINK, J.H. (1992a) Species extinction and concomitant ecological changes in Lake Victoria. *Neth. J. Zool.* **42**, 214–232.
- WITTE, F., GOLDSCHMIDT, T. & WANINK, J.H. (1995) Dynamics of the haplochromine cichlid fauna and other ecological changes in the Mwanza Gulf of Lake Victoria. In: *The Impact of Species Changes in African Lakes* (Eds T. J. PITCHER and P. J. B. HART). Chapman & Hall, London.
- WITTE, E., GOLDSCHMIDT, T., WANINK, J., VAN OIJEN, M., GOUDSWAARD, K., WITTE-MAAS, E. & BOUTON, N. (1992b) The destruction of an endemic species flock: quantitative data on the decline of the haplochromine cichlids of Lake Victoria. *Env. Biol. Fish.* **34**, 1–28.
- WITTE, F. & GOUDSWAARD, P.C. (1985) Aspects of the haplochromine fishery in southern Lake Victoria. In: *Report on the Third Session of the Sub-committee for the Development and Management of the Fisheries of Lake Victoria, 4–5 October 1984, Jinja, Uganda (CIFA)*. FAO Fisheries Report 335. FAO, Rome.

(Manuscript accepted 18 April 2001)