

Short Communication

Length-Weight Relationship and Condition Factor (K constant) of *Dentex maroccanus*, Valenciennes 1830 (Family Sparidae) at Malindi, Kenya

Christopher M. Aura^{1,2}, Rashid O. Anam¹, Safina Musa³ and Edward Kimani¹

¹Kenya Marine and Fisheries Research Institute PO Box 81651-80100, Mombasa, Kenya;

²Laboratory of Marine Bioresource & Environment Sensing, Hokkaido University, Minatocho, Hakodate, Hokkaido 041-8611, Japan; ³Kegati Aquaculture Research Center, Kenya Marine and Fisheries Research Institute, PO Box 3259-40200, Kisii, Kenya.

Keywords: *Dentex maroccanus*, Sparidae, length-weight relationship, condition factor, Kenya

Abstract—Length-weight relationships and condition factors of Sparidae along the Kenyan coast are unknown. This study investigated the length-weight relationship and condition factor of *Dentex maroccanus* Valenciennes, 1830, a sparid, found in Malindi, Kenya, from June to August 2008. The length-weight coefficients were $a = 2.04$ and $b = 2.87$ to 2.96 . The condition factor ranged from 0.91 to 2.84 . Regression coefficients obtained were indicative of positive allometric growth.

INTRODUCTION

Dentex maroccanus (Valenciennes, 1830) is a member of the family Sparidae, Order Perciformes, commonly known as fin-rayed sea breams. Sparids are chiefly marine, being rare in fresh and brackish water (De Innocentiis *et al.*, 2005; Omogoriola *et al.*, 2011). They are widely distributed in the tropical and temperate Atlantic, Indian and Pacific Oceans. Many species have been found to be hermaphroditic, having male and female gonads simultaneously, while others change their sex as they grow. The family is of importance to fisheries based on their food value rather than the abundance of any species in particular (Omogoriola *et al.*, 2011).

Length-weight relationships provide useful information on fish species within a

given geographic region (Morato *et al.*, 2001; Aura *et al.*, 2011). In fish, size is generally more biologically relevant than age, mainly because several ecological and physiological factors are more size-dependent than they are age-dependent. Therefore, variability in size has important implications in fisheries science and population dynamics (Erzini, 1994), and is one of the most common measurements in fisheries data (Mendes *et al.*, 2004). While aspects of the biology of *D. maroccanus* have been investigated by several authors (e.g. Bauchot & Hureau 1986, 1990; Bauchot, 1987), this is the first study of the length-weight relationship of this important fish species which is landed in quantities in Kenya, particularly between May and August during its spawning period.

MATERIALS and METHODS

Study areas

Malindi is a coastal town near the south of Malindi-Ungwana Bay which lies between 3°30'S - 2° 30'S and 40° 00'N - 41° 00'N (Fig. 1). The bay falls within the Malindi and Tana River Districts in the central part of the Coast Province. Fishing and tourism are the main income sources of people in this area (Aura *et al.*, 2011).

Malindi-Ungwana Bay is under the influence of the dominant offshore current regimes. During the South East Monsoon (SEM), which occurs between April and October, the current circulation is dominated by the northward flow of the East African Coastal Current. During this season, the bay

also receives the heaviest inflow of the Tana and Sabaki Rivers. During the North East Monsoon (NEM), between November and March, the northward flowing East African Coastal Current joins the southward flowing Somali current to form the Equatorial Counter Current. In addition, the bay is shallow with a wide continental shelf ranging between 15 and 60 km. The mean depth at high spring tide is 12 m at 1.5 NM offshore and 18 m at 6.0 NM. The depth increases rapidly to 100 m beyond this. Terrigenous sediments from the rivers dominate the substratum in the bay (Aura *et al.*, 2011).

Length-weight data of 688 specimens of *Dentex maroccanus* were obtained from artisanal fishing catches at Malindi landing sites, harvested with fishing gear such as gill nets, seine nets, hand lines and traps. The data were collected daily in June, July and August 2008 for an equal number of males and females. The fish were sexed by macroscopic examination of the gonads according to Grandcourt *et al.* (2004). Standard length (SL) and total length (TL) were measured using a measuring board to the nearest mm. The body weight of the fish was measured to the nearest 0.1 g with a digital balance after removing excess water and adherent matter such as sand. The length-weight data were recorded in a Microsoft Excel spreadsheet for analysis.

The length-weight relationship and condition factor were respectively derived using the equations $\log W = \log a + b \cdot \log L$ and $K = 100W/L^b$ (Ricker, 1973) where: W = weight (g), L = total length (cm), a = regression intercept, b = growth regression slope and K = condition factor. The slopes of the length-weight regressions were compared for individual and combined sexes to determine the allometric growth of *D. marocannus* according to Huxley (1924).




Figure 1. Location of Malindi-Ugwana Bay on the coast of Kenya.




Figure 2. Length-weight relationship of *Dentex maroccanus* males (n = 344).

RESULTS

Table 1 lists parameters of the length-weight relationships of *Dentex macrocannus* (Figs 2-4) measured during the study period. The mean value and range of the condition factor of males was higher than for females and a t-test showed that these differences (p = 0.04) were significant (Table 2).

Most specimens fell within the size classes of 24.1 – 30.0 cm (n = 305) and 18.0 - 24.0 cm (n = 195). Only 30 specimens fell into the largest size class (35.1 - 41.0 cm). Most specimens fell within the 151 - 300 g (n = 250) and 301 - 450 g (n = 198) weight classes but nine fell in the heaviest weight class of 751 - 848 g.

Table 1. Parameters for length-weight relationships of *Dentex maroccanus*.

Sex	TL (cm)	Weight (g)	Mean SL (cm)	a	b
Male	15.7-40.3	39-820	23.1	2.04	2.87
Female	12.6-32.0	41-780	20.3	2.03	2.96
Combined sexes	12.6-40.3	39-820	20.7	2.04	2.91

The allometric growth coefficients for each sex and the sexes combined fell within the expected range of values (b = 2.5 - 3.5) in the literature (Omogoriola *et al*, 2011; <http://www.fishbase.org/summary/Dentex-maroccanus.html>) and compared favourably with other sparids (Table 3). As they were >1, they are indicative of positive allometric growth (Huxley, 1924).



Acknowledgments—Kenya Marine and Fisheries Research Institute in Mombasa, Kenya provided the funds and logistics of this study.

Table 2. Condition factor of *Dentex maroccanus* in June and July 2008 and of both sexes at Malindi.

Month/Sex	Range of K	Mean K
June	0.73-1.07	0.99
July	0.83-2.70	1.02
Male	1.45-2.87	2.49
Female	0.79-2.01	1.17
Combined sexes	0.91-2.84	2.55

Table 3. Length-weight relationships of some Sparidae.

Species name (Linnaeus, 1758)	n	Mean TL \pm SE	log(a)	b	r ²	Author
<i>Sparus aurata</i>	395	19.2 \pm 0.05	0.03	2.67	0.91	Cherif <i>et al.</i> (2008)
<i>Spicara maena</i>	330	15.4 \pm 0.09	0.014	2.87	0.96	Cherif <i>et al.</i> (2008)
<i>Boops boops</i>	243	18.5 \pm 0.14	0.007	3.06	0.98	Cherif <i>et al.</i> (2008)
<i>Dentex congolensis</i>	308	14.7 \pm 2.57	-1.61	2.79	0.91	Omogoriola <i>et al.</i> (2011)
<i>Dentex angolensis</i>	277	16.47 \pm 3.76	-1.56	2.77	0.98	Omogoriola <i>et al.</i> (2011)

Figure 3. Length-weight relationship of *Dentex maroccanus* females (n = 344).Figure 4. Length-weight relationship of *Dentex maroccanus*: combined sexes (n = 668).

References

- Aura M C, Munga CN, Kimani EN, Manyala JO, Musa S (2011) Length-weight relationships for nine deep sea species off the Kenyan coast. *Pan-American Journal of Aquatic Sciences* 6: 188-192
- Ayoade AA, Ikulala AO (2007) Length-weight relationship, condition factor and stomach contents of *Hemichromis bimaculatus*, *Sarotherodon melanotheron* and *Chromidotilapia guentheri* (Perciformes: Cichlidae) in Eleiyele Lake, Southwestern Nigeria. *International Journal of Tropical Biology* 55: 969-97
- Bauchot M-L, Hureau J-C (1990) Sparidae. In Quero JC, Hureau JC, Karrer C, Post A, Saldanha L (eds) Check-list of the fishes of the eastern tropical Atlantic (CLOFETA). JNICT, Lisbon; SEI, Paris; and UNESCO, Paris, Vol. 2, pp 790-812
- Bauchot M-L (1987) Sparidae. In: Fischer W, Bauchot M-L, Schneider M (eds) Fiches FAO d'identification des espèces pour les besoins de la pêche (Révision 1). Méditerranée et mer Noire. Zone de pêche 37. Vol. II, Vértébrés, pp 761-1530
- Bauchot M-L, Hureau J-C (1986) Sparidae. In: Whitehead PJP, Bauchot M-L, Hureau J-C, Nielsen J, Tortonese E (eds) Fishes of the North-eastern Atlantic and the Mediterranean (FNAM). Unesco, Paris, Vol. II, pp 883-907
- Cherif M, Zarrad R, Gharbi H, Missaoui H, Jarboui O (2008) Length-weight relationships for 11 fish species from the Gulf of Tunis (SW Mediterranean Sea, Tunisia). *Pan-American Journal of Aquatic Sciences* 3: 1-5
- De Innocentiis S, Miggiano E, Ungaro A, Livi S, Sola L, Crosetti D (2005) Geographical origin of individual breeders from gilthead sea bream (*Sparus auratus*) hatchery brood stocks inferred by microsatellite profiles. *Aquaculture* 247: 227-232
- Erzini K (1994) An empirical study of variability in length at age of marine fishes. *Journal of Applied Ichthyology* 10: 17-41
- Grandcourt EM, Al Abdessalaam TZ, Francis F, Al Shamsi AT (2004) Biology and stock assessment of the sparids, *Acanthopagrus bifasciatus* and *Argyrops spinifer* (Forsskål, 1775), in the Southern Arabian Gulf. *Fisheries Research* 69: 7-20
- Huxley J (1942) Constant differential growth-ratios and their significance. *Nature* London 114: 895-1932
- Jones RE, Petrell RJ, Paulty D (1999) Using modified length-weight relationships to assess the condition of fishes. *Aquacultural Engineering* 20: 261-276
- Mendes B, Fonseca P, Campos A (2004) Weight-length relationships for 46 fish species of the Portuguese west coast. *Journal of Applied Ichthyology* 20: 355-361
- Morato T, Afonso P, Loirinho P, Barreiros JP, Sanstos RS, Nash RDM (2001) Length-weight relationships for 21 coastal fish species of the Azores, North-eastern Atlantic. *Fisheries Research* 50: 297-302
- Omogoriola HO, Williams AB, Majisola OA, Olakolu FC, Ukaonu FU, Myade EF (2011) Length-weight relationships, condition factor (K) and relative condition factor (Kn) of sparids, *Dentex congoensis* (Maul, 1954) and *Dentex angolensis* (Maul and Poll, 1953), in Nigerian coastal water. *International Journal of Biological and Chemical Sciences* 5: 739-747
- Ricker WE (1973) Linear regressions in fishery research. *Journal of Fisheries Research Board of Canada* 30: 409-434