




Detection of tilapia lake virus (TiLV) infection by PCR in farmed and wild Nile tilapia (*Oreochromis niloticus*) from Lake Victoria

K K Mugimba^{1,2*} | A A Chengula^{1,3*} | S Wamala^{1,2} | E D Mwege^{1,3} | C J Kasanga³ |
D K Byarugaba² | R H Mdegela³ | S Tal⁴ | B Bornstein⁴ | A Dishon⁴ | S Mutoloki¹  |
L David⁵  | Ø Evensen¹ | H M Munang'andu¹ 

¹Faculty of Veterinary Medicine and Biosciences, Department of Basic Sciences and Aquatic Medicine, Section of Aquatic Medicine and Nutrition, Norwegian University of Life Sciences, Oslo, Norway

²Department of Biotechnical and Diagnostic Sciences, College of Veterinary Medicine Animal Resources and Biosecurity, Makerere University, Kampala, Uganda

³Department of Microbiology, Parasitology and Immunology, College of Veterinary Medicine and Biomedical Sciences, Sokoine University of Agriculture, Morogoro, Tanzania

⁴KoVaX Ltd., Jerusalem, Israel

⁵R.H. Smith Faculty of Agriculture, Food and Environment, Department of Animal Sciences, The Hebrew University of Jerusalem, Rehovot, Israel

Correspondence

Hetron Mweemba Munang'andu, Faculty of Veterinary Medicine and Biosciences, Department of Basic Sciences and Aquatic Medicine, Section of Aquatic Medicine and Nutrition, Norwegian University of Life Sciences, Oslo, Norway.
Email: hetryonej.
mweemba.munangandu@nmbu.no

Funding information

NORAD TRAHESA, Grant/Award Number: TAN/13/0027

Abstract

Tilapia lake virus disease (TiLVD) has emerged to be an important viral disease of farmed Nile tilapia (*Oreochromis niloticus*) having the potential to impede expansion of aquaculture production. There is a need for rapid diagnostic tools to identify infected fish to limit the spread in individual farms. We report the first detection of TiLV infection by PCR in farmed and wild Nile tilapia from Lake Victoria. There was no difference in prevalence between farmed and wild fish samples ($p = .65$), and of the 442 samples examined from 191 fish, 28 were positive for TiLV by PCR. In terms of tissue distribution, the head kidney (7.69%, $N = 65$) and spleen (10.99%, $N = 191$), samples had the highest prevalence ($p < .0028$) followed by heart samples (3.45%, $N = 29$). Conversely, the prevalence was low in the liver (0.71%, $N = 140$) and absent in brain samples (0.0%, $N = 17$), which have previously been shown to be target organs during acute infections. Phylogenetic analysis showed homology between our sequences and those from recent outbreaks in Israel and Thailand. Given that these findings were based on nucleic acid detection by PCR, future studies should seek to isolate the virus from fish in Lake Victoria and show its ability to cause disease and virulence in susceptible fish.

KEYWORDS

Lake Victoria, Nile tilapia, PCR, phylogenetic, surveillance, tilapia lake virus

1 | INTRODUCTION

Tilapia lake virus (TiLV), also known as syncytial hepatitis of tilapia—SHT, was first identified and shown to cause mortalities in Nile tilapia (*Oreochromis niloticus*) in 2012 in Israel by KoVax, (personal communication), following summer mortalities in tilapia fish farms in

Israel. Soon after it was reported that the same virus was present in tilapia fish in the Sea of Galilee in Israel and again that this virus was causing disease and mortalities in Nile tilapia (Eyngor et al., 2014). It has since been associated with outbreaks in Colombia, Ecuador, Egypt, Israel and Thailand (Bacharach et al., 2016; Del-Pozo et al., 2017; Fathi et al., 2017; Kembou Tsofack et al., 2017; Nicholson et al., 2017; Surachetpong et al., 2017). Based on motif alignment of its segment-1 with the PB1 segment of influenza A, B and C, the

*Equal contribution

aetiological agent has been characterized as an orthomyxo-like virus (Bacharach et al., 2016). Classification by the International Committee of Virus Taxonomy (ICVT) puts TiLV as a single new species known as *Tilapia tilapinevirus* in the new genus *Tilapinevirus* (Adams et al., 2017). It is made of 10 segments unlike other orthomyxoviruses such as influenza that are made up of eight segments (Palese & Schulman, 1976). The length of the total viral genome is about 10,323 kp (Bacharach et al., 2016, Bacharach et al., 2016; Del-Pozo et al., 2017, Eyngor et al., 2014). In vitro studies show that the virus grows well at 23–30°C in vitro. Studies carried out this far show that mortalities occur at temperatures above 25°C as shown from the summer die-offs associated with TiLV in Egypt and Israel (Fathi et al., 2017). TiLV has so far been reported in Nile tilapia (Egypt, Thailand) (Fathi et al., 2017; Nicholson et al., 2017; Surachetpong et al., 2017), red tilapia (Thailand) (Surachetpong et al., 2017) and the hybrid tilapia *O. niloticus* × *O. aureus* (Israel) (Bacharach et al., 2016), suggesting that the range of the tilapines susceptible to TiLV could be wider.

Lake Victoria is the world's second largest freshwater lake covering a surface area of 68,000 km² shared by three countries in East Africa namely Kenya (6%), Uganda (45%) and Tanzania (49%). By the 1960s, it was habitat for several fish species dominated by the tilapine cichlids such as *O. esculantus* and *O. variabilis* and home to more than 200 haplochromine cichlids (Goudswaard & Witte, 1997; Goudswaard, Witte, & Katunzi, 2002; Kudhongania & Cordone, 1974; Ogutu-Ohwayo, 1990). Nile tilapia and Nile perch (*Lates niloticus*) were introduced in the 1950s to replace the declining tilapine species, which led to disappearance of >50% of the indigenous fish species in Lake Victoria (Ogutu-Ohwayo, 1990). Since then, Nile tilapia and Nile perch species continued to increase although by 2002, the Nile perch population began to decline giving way for the Nile tilapia to become the most dominant fish species in Lake Victoria (Njiru et al., 2012; Ogutu-Ohwayo, 1994; Witte et al., 1991). The recent introduction of cage farming of Nile tilapia further increases its dominance on Lake Victoria. The emergence of viral diseases such as Tilapia lake virus disease (TiLVD) poses a significant threat to the expansion of Nile tilapia production, which has tremendously increased to become one of the leading cultured fish species in the world in the last decade. Furthermore, the rapid rate at which the disease is being reported to cause outbreaks in different continents across the world (Fathi et al., 2017; Nicholson et al., 2017; Surachetpong et al., 2017) calls for the development of rapid diagnostic tools for prompt virus identification to pave way for the design of timely disease control strategies. Thus far, diagnosis of reported outbreaks has mainly been based on virus isolation, characterization, culture followed by reinfection in susceptible fish to demonstrate the characteristic syncytial hepatitis and other pathological lesions in susceptible fish (Del-Pozo et al., 2017; Tattiyapong, Dachavichitlead, & Surachetpong, 2017; Tsofack et al., 2016). Although these steps fulfil the Koch's postulates by establishing the disease-causal factor relationship, virus isolation, culture and reinfection are not ideal for surveillance programmes especially in situations with high number of samples. This is because the culture and reinfection approach is not only expensive, but could take long to generate results. Hence, there

is urgent need for rapid diagnostic tests suitable for surveillance programmes to expedite the process of establishing the distribution of TiLV. Moreover, developing surveillance diagnostic tools would pave way to designing appropriate disease control measures aimed at preventing the spread of the virus in the aquaculture industry. The aim of this study was twofold: (i) to develop and optimize a PCR-based method for the detection of TiLV and (ii) to investigate the possible existence of TiLV in Nile tilapia found in Lake Victoria.

2 | METHODS AND MATERIALS

2.1 | Sample collection and study sites

Nile tilapia samples were collected from the Ugandan and Tanzanian parts of Lake Victoria in 2016 and 2015, respectively. For Ugandan samples, 83 fish were collected from 14 sampling sites and transported to Makerere University on ice in cool boxes. Among these, seven sites were from cage-farmed fish while the other seven were from wild fish (Table 1). Sampling for the wild sites was performed in areas at least 20–50 km into the Lake away from the shore with minimum distances of 20 km apart while the cage farms were within 10 km from the lakeshore. Fish were dissected and processed at the Faculty of Veterinary Medicine of Makerere University. All tissues collected were stored in RNAlater and stored at 4°C for 24 hr followed by –80°C until transfer to the Norwegian University of Life Sciences (NMBU) in Oslo, Norway. In Tanzania, a total of 216 samples were collected from 108 wild fish at four sampling sites (Table 1). Two sampling sites approximately 20 km apart were sampled in Maganga beach area and another two sites in the Mchongmani area separated by a distance of approximately 25 km apart. Dissections were carried out at the Fisheries Education and Training Authority (FETA) laboratory in Mwanza, and the tissues collected were stored in RNAlater at –20°C for 5 days. Thereafter, all samples were transported to the College of Veterinary Medicine and Biomedical Sciences at Sokoine University of Agriculture in Morogoro where they were stored at –80°C until shipment to NMBU. Overall, a total of 442 organs from 191 fish were collected from Lake Victoria as summarized in Table 1.

2.2 | Virus propagation and cell culture

Tilapia cell cultures were generated from hybrid *Oreochromis niloticus* × *Oreochromis aureus*. In brief, caudal fins were removed from killed fish that weighed 30 g. Fish were then bathed in 1% sodium hypochlorite solution for 1 min and then rinsed in 70% ethyl alcohol. Fins were washed three times in phosphate buffer saline (PBS) containing 10% penicillin–streptomycin and 2.5% nystatin. The fins were transferred to Petri dishes, extensively minced with scissors, and semi-dry small tissue pieces of approximately 1 mm³ were placed in dry 50-ml culture flasks (Nunc, Denmark). After 24-hr incubation at room temperature, the clumps adhering to the flasks were covered with Leibovitz (L-15) medium (Sigma) supplemented with 10% FBS (Biological Industries, Israel), 1% nystatin and 2.5%

TABLE 1 Sampling sites and number of fish samples

Country	Sampling site	Culture system	Positive fish/total	Positive samples/total	Organs (Positive/total)				
					Liver	Heart	Head kidney	Spleen	Brain
Tanzania	Maganga beach-1	Wild	4/19	4/38	0/19	–	–	4/19	–
	Mchongomani-1		7/28	7/56	0/28	–	–	7/28	–
	Maganga beach-2		6/35	6/70	0/35	–	–	6/35	–
	Mchongomani-2		1/26	1/52	0/26	–	–	1/26	–
Uganda	Kigungu	Cage farms	5/8	5/32	0/8	0/8	3/8	2/8	–
	Lwera		1/8	1/16	–	–	1/8	0/8	–
	Kasenyi		0/8	0/8	–	–	–	0/8	–
	Entebbe		0/5	0/5	–	–	–	0/5	–
	Bukanama		0/5	0/5	–	–	–	0/5	–
	SON		1/6	1/23	1/6	0/6	0/5	0/6	–
	Kome		1/5	1/11	–	–	1/6	0/5	–
	Lake Victoria-1	Wild	0/5	0/10	–	–	0/5	0/5	–
	Lake Victoria-2		0/6	0/12	–	–	0/6	0/6	–
	Lake Victoria-3		0/6	0/12	–	–	0/6	0/6	–
	Lake Victoria-4		0/3	0/6	–	–	0/3	0/3	–
	Lake Victoria-5		1/6	1/30	0/6	0/6	0/6	1/6	0/6
	Lake Victoria-6		1/6	1/30	0/6	1/6	0/6	0/6	0/6
	Lake Victoria-7		0/6	0/26	0/6	0/3	0/6	0/6	0/5
Total			28/191	28/442	1/140	1/29	5/65	21/191	0/17
Prevalence			14.66%	6.33%	0.71%	3.45%	7.69%	10.99%	0.00

penicillin–streptomycin. Cells were maintained at 28°C in a CO₂-free environment. At 10- to 14-day incubation, cells grew out from the tissue to form a monolayer around each clump. The monolayer cultures were trypsinized and transferred into new flasks with fresh medium. The cells have been passed for over 100 times to form a stable cell line and are referred to as tilapia fin cells—TFC#10.

The virus used as a positive control in this study was provided by KoVax Vaccine Company in Israel. Virus isolation from sick fish was performed as follows: sick fish showing signs of apathy, reduced appetite and mortality were collected and frozen at –80°C. Kidney, spleen, intestine, gills and brains were collected and homogenized in PBS. The homogenate was filtered through a 0.2-µm filter (Sartorius). Filtered homogenates were used to inoculate naïve TFC#10 cultures, incubated at 28°C and monitored daily. Cytopathic effect (CPE) appeared at 4–7 days post-inoculation. Once extensive CPE was evident, virus suspension was harvested, aliquoted and stored at –80°C for further use.

For the negative control cells, PBS only was used instead of the virus for adsorption. After 7 days of incubation, suspensions for both virus-infected and virus-non-infected cells were harvested and used for RNA extraction as described below.

2.3 | RNA extraction and cDNA synthesis

Extraction of total RNA from the 442 samples (Table 1) was carried out using a combination of the TRIzol® (GIBCO, Life Technologies)

and RNeasy Mini kit (Qiagen, Hilden, Germany) techniques as previously described (Munang'andu, Fredriksen, Mutoloki, Dalmo, & Even- sen, 2013; Munang'andu et al., 2012; Munang'andu, Sandtrø, et al., 2013). Briefly, approximately 30 mg of tissue was homogenized in 1 mL TRIzol followed by centrifugation at 12,000 g for 10 min at 4°C. Thereafter, the supernatant was transferred into an Eppendorf tube followed by addition of 0.2 ml chloroform to each sample. After vortexing for 15s, samples were left for 5 min at room temperature followed by spinning at 12,000 g for 15 min. The aqueous phase was transferred into another Eppendorf tube. After adding 0.6 ml of 70% ethanol, the tubes were vortexed and the contents were transferred to RNeasy spin columns. Thereafter, the Qiagen protocol was used based on the manufacturer's guidelines (Qiagen, Hilden, Germany). RNA quantification was carried out using a spectrophotometer (NanoDrop® ND-1000, Thermo Scientific Inc). The synthesis of cDNA was carried out in 20 µl reaction volumes using the Transcrip- tor First Strand cDNA Synthesis Kit that has an integrated step for the removal of contaminated genomic DNA (Qiagen). The final cDNA was stored at –80°C until use.

Preparation of the negative control samples was made by extract- ing RNA and cDNA synthesis from the non-infected TFC#10 cells while RNA and cDNA synthesized from infected cells were used to prepare the virus-positive controls. In addition, a second negative control was prepared by extracting RNA from head kidney, spleen and liver samples collected from six fish of the 15th generation of Nile tilapia cultured by the GIFT project cultured at the NMBU followed by

TABLE 2 Primer sequences

Segment	Primer sequence	Length (bp)	T _m (°C)
Segment-1	FWD-CCTCATTCTCGTTGTGTAAGT	1,000	62
	REV-AGGAGTTGCTGTTGGGTATAG		
Segment-2	FWD-GTCCAGGGCGGTATGTATTG	834	62
	REV-CTTACGGCTGACAAGTCTCTAAG		
Segment-3	FWD-GTCGAGGCATTCCAGAAGTAAG	834	62
	REV-GAGCTAAGGGAACGGCTATTG		
Segment-4	FWD-GCCTACTTCGTTGCCTATCTC	524	62
	REV-GCCCAATGGTCCCATATCT		
Segment-5	FWD-CAACTCTTAGCCTCCGGAATAC	696	62
	REV-CGTTCTGCACTGGGTTACA		
Segment-6	FWD-CCCACACGACAGGACATATAG	948	62
	REV-GAGTTGGCTTAGGGTGATAAGA		
Segment-7	FWD-TCCTTTAGGGATTGCACTAAC	486	62
	REV-TTCCATCGACTGCTCCTAGA		
Segment-8	FWD-CTTAAGGGCCATCCTGTCATC	476	62
	REV-TGGCTCAAATCCCAACTAA		
Segment-9	FWD-GATATCCTCCACATGACCCTTC	261	62
	REV-GTACGTCACCTTTGTGCCATTAC		
Segment-10	FWD-TCCTCTCTGTCCCTTCTGTT	276	62
	REV-CAGGATGAGTGTGGCAGATTAT		

cDNA synthesis. The cDNA prepared from the GIFT fish samples was pooled for use as negative control from a population not previously exposed to TiLV. Hence, the negative control samples used in this study were designated as TCF#10 cells and GIFT tissue.

2.4 | Optimization of the polymerase chain reaction test

A total of 10 primers (Table 2) were designed targeting the 10 segments of the TiLV genome. For PCR optimization, each primer pair was tested against two TiLV positive controls designated as 2V and 5V, two GIFT tissue negative controls designated as 4T and 5T, two TFC#10 cells negative control, namely 2C and 3C, and one sterile water negative control (NC). The objective of using two replicates for each control sample was to compare the reproducibility of the PCR products generated after amplification between duplicates. Further, the purpose of using two negative controls (TFC#10 cells and pooled GIFT tissue cDNA) was to compare the reliability of a continuous cell line and host tissue-derived negative control during PCR optimization. Hence, each primer was tested against a total of seven samples in order to identify primers that only detect viral cDNA in order to reduce the chances of producing unspecific PCR products. All PCRs for amplification of the segment 1–10 genes were carried out using the Q5 High-Fidelity DNA Polymerase (New England BioLabs, Inc.). After gel electrophoresis analyses, only primers showing bands in the TiLV positive controls without bands in the TFC#10 cells and GIFT tissue negative controls were selected for use in the screening of Nile tilapia samples for the presence of TiLV in the next step.

2.5 | Screening and sequencing of Nile tilapia samples from Lake Victoria

Once the PCR optimization process was completed, the selected primers were used to screen Nile tilapia samples from Lake Victoria for the presence of TiLV. As shown in Table 1, a total of 442 samples were examined from different organs including heart, liver, brain, head kidney and spleen. All the 28 PCR products obtained from the screening of Nile tilapia samples shown in Table 1 were extracted and purified using the QIAquick Gel extraction kit according to manufacturer's instruction (Qiagen, Hilden, Germany). Amplification of the TiLV segment-2 genes was performed using the Q5 High-Fidelity DNA Polymerase as described above. PCR products were then separated using 1.5% agarose gel electrophoresis and extracted using the QIAquick Gel Extraction Kit (Qiagen, Hilden, Germany). Sequencing was performed on a commercial basis by GATC Biotech (<https://www.gatc-biotech.com>). The CLC Workbench 6.0 (www.clcbio.com) and Mega7 software (Kumar, Stecher, & Tamura, 2016) were used for sequence alignment and phylogenetic tree analyses. Phylogenetic trees were inferred by the maximum-likelihood method, bootstrapped 1,000 times based on the JTT+G matrix-based model (Jones, Taylor, & Thornton, 1992). Initial tree(s) for the heuristic search were obtained automatically by applying neighbour-joining and BioNJ algorithms to a matrix of pairwise distances estimated using the maximum composite likelihood (MCL) approach, and then selecting the topology with superior log-likelihood value. The tree was drawn to scale, with branch lengths measured in the number of substitutions per site. The analysis involved 36 nucleotide

sequences. Codon positions included were 1st+2nd+3rd+non-coding. All positions containing gaps and missing data were eliminated. There were a total of 270 positions in the final data set. Evolutionary analyses were conducted in MEGA7 (Kumar et al., 2016). The results obtained for the different groups and organs were analysed statistically using Fisher's exact test using GraphPad Prism version 7.00 for Windows, GraphPad Software, La Jolla, California, USA.

3 | RESULTS

3.1 | Polymerase chain reaction optimization

Figure 1 shows that PCR products obtained for segments 1, 3, 6, 8, 9 and 10 had strong bands in the TiLV positive controls (lanes 2V and 5V) and faint bands in the GIFT tissue (lanes 4T and 5T) and no bands were detected in TFC#10 cells (lanes 2C and 3C) and sterile water (lane N). Although the presence of strong bands in the virus-positive controls show that these primers detected the viral cDNA, the presence of faint bands in the GIFT tissue negative controls shows that they produced unspecific amplifications. In addition, there were no PCR products for segment-4 for the positive and negative controls while PCR products for segment-7 had weak bands in the virus control and no bands were seen in the negative controls (Figure 1). However, PCR products for segment-2 were only detected in the virus controls (2V and 5V) and no bands were detected in the GIFT fish (lanes 4T and 5T), TFC#10 cells (lanes 2C and 3C) and sterile water (lane N) negative controls. Therefore, the presence of clear bands in the TiLV positive controls (lanes 2V and

5V) and the absence of PCR products in the negative controls were indicative that segment-2 primers were only able to detect TiLV cDNA, but not unspecified amplifications. Therefore, segment-2 primers were selected for the screening of Nile tilapia samples from Lake Victoria in the next step based on their ability to only detect viral cDNA and not host DNA, while primers for other segments were considered less suitable because either they gave some levels of unspecific amplifications in the controls or failed to detect the viral cDNA. Finally, the GIFT fish negative control was more reliable at detecting unspecific amplifications (Figure 1, lanes 4T and 5T) compared to the TFC#10 cells negative control (lanes 2C and 3C) that showed absence of unspecific amplifications for primers tested during the PCR optimization process.

3.2 | Screening of Nile tilapia samples from Lake Victoria

Table 1 shows a summary of the number of samples examined for the presence of TiLV nucleic acids by PCR using segment-2 primers. Of the 191 fish examined, 28 were found positive by PCR for TiLV nucleic acids with a prevalence of 14.66% ($N = 191$). The prevalence in caged and wild fish was 17.78% ($N = 45$) and 13.70% ($N = 146$), respectively. There was no significant difference in the prevalence ($p = .136$) among wild fish from the Tanzania side (16.67%, $N = 108$) compared with the Uganda side (5.30% $N = 38$). However, there was a significant difference ($p < .0028$) in tissue distribution among organs examined. PCR products were detected in 10.99% ($N = 191$) spleen, 7.69% ($N = 65$) head kidney, 3.45% ($N = 29$) heart and liver




FIGURE 1 Shows electrophoresis gel analysis of TiLV positive control and the TFC#10 cells and GIFT fish negative control samples tested against TiLV segment 1–10 for the 10 primers (P1–P10) enlisted in Table 1. Note that both the positive and negative control samples are tested in duplicates in which the TFC#10 cells cDNA samples are designated as 2C and 3C, GIFT fish cDNA samples are designated as 4T and 5T while the positive virus (TiLV) control samples are designated as 2V and 5V. In addition, a single lane designated as NC for RNase-free water was added to each segment tested. The expected amplicon for each primer is shown alongside the name of the segment tested. There were no detectable bands in the TFC#10 cell (lanes 2C and 3C) and RNase-free water (lane NC) negative controls for all the 10 primers tested for segments 1–10. Note that P1, 2, 3, 6, 8, 9 and 10 showed clear bright bands of the virus-positive control (lanes 2V and 5V) while P7 had a faint band in lane 2V and no bands were detected in P4. Finally, the GIFT fish samples showed faint bands in P1, 3, 6, 8, 9 and 10 in lanes 4T and 5T of variable amplicon sizes

0.71% ($N = 140$) samples while no PCR products were detected in brain samples (0.0%, $N = 17$). In summary, Table 2 shows that the lymphoid organs, mainly comprising of the head kidney and spleen, had the highest prevalence followed by heart samples.

3.3 | Sequence alignment and phylogenetic analysis

Table 3 provides a summary of sequenced samples showing their origin, organ, size of the sequence product and GenBank accession numbers. The average length of the 28 sequences retrieved was 768 bp. A blast analysis against the NCBI sequence database showed that all sequences obtained were homologous to TiLV segment-2 sequences obtained from viruses isolated from Israel and Thailand. The samples clustered into two groups that were slightly different from each other and corresponded to different isolates that were already reported in the GenBank database (Figure 2). Group I comprised of a total 25 Lake Victoria sequences of which nine were from Uganda and 16 from Tanzania that were identical with the Israeli KU552132 sequence deposited by Tal et al., (2016) and

Thailand KX631922.1 sequence deposited by Surachetpong et al., (2017). Group II consisted of only three Lake Victoria sequences of which one was from Uganda and two were from Tanzania that were closely related to the Israeli KU751815.1 and NC029921 sequences deposited by Eyngor et al., (2014) and (Bacharach et al., (2016) , respectively.

4 | DISCUSSION

In this study, we detected TiLV nucleic acids in an area with no record of previous outbreaks. In line with Louws, Rademaker, & de Bruijn, (1999) who pointed out that the three Ds of PCR analyses are detection, diversity and diagnosis, we have (i) detected, (ii) shown phylogenetic diversity and (iii) diagnosed the presence of TiLV nucleic acids in Nile tilapia from Lake Victoria using PCR. Although we did not fulfil the Koch's postulate (Evans, 1976; Fredericks & Relman, 1996; Gradmann, 2014) by establishing the disease-casual factor relationship based on isolation, characterization, culture and

TABLE 3 Description of samples used for TiLV sequencing

Sample ID	Country	Source	Organ	Size (bp)	GenBank Acc #
UG2016-01	Uganda	Wild	Head kidney	830	MF536423
UG2016-02	Uganda	Wild	Liver	782	MF536429
UG2016-03	Uganda	Wild	Head kidney	332	MF536432
UG2016-04	Uganda	Wild	Heart	805	MF536426
UG2016-05	Uganda	Wild	Spleen	818	MF536427
UG2016-06	Uganda	Wild	Head kidney	830	MF536424
UG2016-07	Uganda	Wild	Spleen	819	MF536425
UG2016-08	Uganda	Wild	Spleen	787	MF536428
UG2016-09	Uganda	Wild	Head kidney	540	MF536430
UG2016-10	Uganda	Wild	Head kidney	724	MF536431
TZ2015-01	Tanzania	Maganga beach	Spleen	777	MF526992
TZ2015-02	Tanzania	Mchongomani	Spleen	767	MF526988
TZ2015-03	Tanzania	Mchongomani	Spleen	827	MF526987
TZ2015-04	Tanzania	Maganga beach	Spleen	755	MF526989
TZ2015-05	Tanzania	Mchongomani	Spleen	828	MF526980
TZ2015-06	Tanzania	Maganga beach	Spleen	669	MF526982
TZ2015-07	Tanzania	Maganga beach	Spleen	706	MF526991
TZ2015-08	Tanzania	Maganga beach	Spleen	675	MF526981
TZ2015-09	Tanzania	Mchongomani	Spleen	725	MF526993
TZ2015-10	Tanzania	Mchongomani	Spleen	827	MF526983
TZ2015-11	Tanzania	Mchongomani	Spleen	578	MF526994
TZ2015-12	Tanzania	Mchongomani	Spleen	758	MF526984
TZ2015-13	Tanzania	Mchongomani	Spleen	792	MF526985
TZ2015-14	Tanzania	Maganga beach	Spleen	576	MF526995
TZ2015-15	Tanzania	Maganga beach	Spleen	731	MF526996
TZ2015-16	Tanzania	Maganga beach	Spleen	765	MF526990
TZ2015-17	Tanzania	Maganga beach	Spleen	794	MF526986
TZ2015-18	Tanzania	Maganga beach	Spleen		




FIGURE 2 Phylogenetic analysis of the 28 Nile tilapia samples from Lake Victoria (TZ for Tanzanian and UG for Ugandan samples) sequenced using segment-2 primers. The evolutionary history was inferred using the maximum-likelihood method based on the Tamura–Nei model (Tamura & Nei, 1993)

reinfection to demonstrate the induction of clinical disease in susceptible fish, our data are strongly suggestive of TiLV infecting Nile tilapia in Lake Victoria. These findings have significant implications on tilapia farming and in countries using parent stocks from Lake Victoria. It is likely that as the demand for high production outputs increases, the use of intensive farming systems based on high stocking densities and artificial feeds aimed at enhancing growth rate is also bound to increase. These factors have the propensity to induce stress in cultured fish (Munang'andu, Mutoloki, & Evensen, 2016), which could lead to underlying viruses to replicate culminating in disease outbreaks. Moreover, high stocking densities are proponents of a high transmission index (Munang'andu et al., 2016), which could increase the risk of inducing TiLV outbreaks in farmed Nile tilapia. Therefore, the detected TiLV nucleic acids in this study serve as an early warning system in which future outbreaks should be thoroughly investigated to confirm the possible existence of TiLV in Nile tilapia in Lake Victoria.

Our findings show an overall population prevalence of 14.66% ($N = 191$), suggesting that in subclinical infection, TiLV could be present in a population at low prevalence only increasing to high levels reaching up to 90% mortality during massive die-offs (Fathi et al., 2017; Surachetpong et al., 2017). Detection of TiLV nucleic acids in the liver, heart, head kidney and spleen in this study is in line with previous studies in which it was shown that TiLV has a tropism for different organs inclusive of the liver, brain, spleen and head kidney when clinical signs of disease are observed (Bacharach et al., 2016; Fathi et al., 2017; Ferguson et al., 2014; Surachetpong et al., 2017; Tsofack et al., 2016). Previous studies have pointed to the brain and liver as target organs of which it has been associated with syncytial formation in the liver as a pathognomonic feature of the disease, at least in one study (Del-Pozo et al., 2017). The high prevalence of TiLV nucleic acids in the head kidney and spleen coupled with a low presence in the liver and absence in brain samples shown in this study suggests that lymphoid organs could be ideal for screening the presence of TiLV nucleic acids during surveillance. However, there is a need for more studies to consolidate this observation.

Phylogenetic analysis clustered our sequences in two similar groups. It is interesting to note that based on segment-2 fragments, these groups correspond with Israeli isolates as shown that group I sequences were clustered together with the Israeli KU552132 sequence while group II sequences were clustered with Israeli KU751815 and NCO29921 sequences, suggesting that TiLV sequences found in Lake Victoria are similar to strains found in the Sea of Galilee in Israel. In addition, group I sequences that formed the largest cluster were similar to the Thailand isolate KX631922, suggesting that TiLV sequences in Thailand, Israel and Lake Victoria might have a common origin. Given that Nile tilapia is originally a freshwater teleost species native to the Nilo-Sudanian ecoregion of Africa (McAndrew, 2000), which in recent decades has been introduced into more than 85 countries in the world (Casal, 2006; Dong, Ataguba, Khunrae, Rattanaojpong, & Senapin, 2017; Molnar, Gamboa, Revenga, & Spalding, 2008), it is likely that its dispersal could have contributed to the spread of TiLV. The existence of TiLV sequences shown in our findings suggests that the virus could have been in existence for a long time such that as tilapia were being dispersed across the world, they carried the virus unnoticed. Its emergence as a fish pathogen is most likely due to stress-related factors induced by current intensified aquaculture systems as well as the increasing environment changes that stress fish in natural waterbodies. However, there is need for detailed studies to determine its distribution and to identify factors linked to its dispersal in aquaculture. Moreover, future studies should seek to establish whether genomic differences seen between group I and II strains in this study account for differences in virulence and persistence linked to subclinical infection, tissue tropism or other factors. Although the homology between sequences obtained in this study and those from previous outbreaks in Israel and Thailand suggests that Lake Victoria sequences could be originating from a virus having the potential to cause outbreaks in Nile tilapia, it is important that these findings are supported with virus isolation, culture and reinfection in future studies.

In summary, this is the first documentation of TiLV genomes in a none-outbreak area (Lake Victoria). The findings clearly demonstrate that viral nucleic acids are present at low level in seemingly healthy fish. Future studies should focus on isolating the virus from Nile tilapia and demonstrate its ability to cause disease in susceptible fish.

ACKNOWLEDGEMENTS

This study was supported in part by the TRAHESA project, which is the Capacity building for training and research in aquatic and environmental health in Eastern and Southern Africa, funded by the Norwegian Development Agency, Project No. TAN/13/0027.

CONFLICT OF INTEREST

Authors declare no competing interests.

ORCID

S Mutoloki  <http://orcid.org/0000-0001-6125-9786>

L David  <http://orcid.org/0000-0002-6555-0594>

H M Munang'andu  <http://orcid.org/0000-0002-1021-6396>

REFERENCES

- Adams, M. J., Lefkowitz, E. J., King, A. M., Harrach, B., Harrison, R. L., Knowles, N. J., ... Mushegian, A. R. (2017). Changes to taxonomy and the international code of virus classification and nomenclature ratified by the international committee on taxonomy of viruses (2017). *Archives of Virology*, *162*, 2505–2538. <https://doi.org/10.1007/s00705-017-3358-5>
- Bacharach, E., Mishra, N., Briese, T., Zody, M. C., Kembou Tsofack, J. E., Zamostiano, R., ... Lipkin, W. I. (2016). Characterization of a novel orthomyxo-like virus causing mass die-offs of tilapia. *MBio*, *7*, e00431–16.
- Casal, C. M. V. (2006). Global documentation of fish introductions: The growing crisis and recommendations for action. *Biological Invasions*, *8*, 3–11. <https://doi.org/10.1007/s10530-005-0231-3>
- Del-Pozo, J., Mishra, N., Kabuusu, R., Cheetham, S., Eldar, A., Bacharach, E., ... Ferguson, H. (2017). Syncytial hepatitis of tilapia (*Oreochromis niloticus* L.) is associated with orthomyxovirus-like virions in hepatocytes. *Veterinary Pathology*, *54*, 164–170. <https://doi.org/10.1177/0300985816658100>
- Dong, H. T., Ataguba, G. A., Khunrae, P., Rattanarojpong, T., & Senapin, S. (2017). Evidence of TiLV infection in tilapia hatcheries from 2012 to 2017 reveals probable global spread of the disease. *Aquaculture*, *479*, 579–583. <https://doi.org/10.1016/j.aquaculture.2017.06.035>
- Evans, A. S. (1976). Causation and disease: The Henle-Koch postulates revisited. *The Yale Journal of Biology and Medicine*, *49*, 175.
- Eyngor, M., Zamostiano, R., Kembou Tsofack, J. E., Berkowitz, A., Bercovier, H., Tinman, S., ... Eldar, A. (2014). Identification of a novel RNA virus lethal to tilapia. *Journal of Clinical Microbiology*, *52*, 4137–4146. <https://doi.org/10.1128/JCM.00827-14>
- Fathi, M., Dickson, C., Dickson, M., Leschen, W., Baily, J., Muir, F., ... Weidmann, M. (2017). Identification of Tilapia Lake Virus in Egypt in Nile tilapia affected by 'summer mortality' syndrome. *Aquaculture*, *473*, 430–432. <https://doi.org/10.1016/j.aquaculture.2017.03.014>
- Ferguson, H., Kabuusu, R., Beltran, S., Reyes, E., Lince, J., & del Pozo, J. (2014). Syncytial hepatitis of farmed tilapia, *Oreochromis niloticus* (L.): A case report. *Journal of Fish Diseases*, *37*, 583–589. <https://doi.org/10.1111/jfd.12142>
- Fredericks, D., & Relman, D. A. (1996). Sequence-based identification of microbial pathogens: A reconsideration of Koch's postulates. *Clinical Microbiology Reviews*, *9*, 18–33.
- Goudswaard, K. P., & Witte, F. (1997). The catfish fauna of Lake Victoria after the Nile perch upsurge. *Environmental Biology of Fishes*, *49*, 21–43. <https://doi.org/10.1023/A:1007311708377>
- Goudswaard, P., Witte, F., & Katunzi, E. (2002). The tilapiine fish stock of Lake Victoria before and after the Nile perch upsurge. *Journal of Fish Biology*, *60*, 838–856. <https://doi.org/10.1111/j.1095-8649.2002.tb02413.x>
- Gradmann, C. (2014). A spirit of scientific rigour: Koch's postulates in twentieth-century medicine. *Microbes and Infection*, *16*, 885–892. <https://doi.org/10.1016/j.micinf.2014.08.012>
- Jones, D. T., Taylor, W. R., & Thornton, J. M. (1992). The rapid generation of mutation data matrices from protein sequences. *Computer Applications in the Biosciences: CABIOS*, *8*, 275–282.
- Kembou Tsofack, J. E., Zamostiano, R., Watted, S., Berkowitz, A., Rosenbluth, E., Mishra, N., ... Bacharach, E. (2017). Detection of Tilapia Lake Virus in clinical samples by culturing and nested reverse transcription-PCR. *Journal of Clinical Microbiology*, *55*, 759–767. <https://doi.org/10.1128/JCM.01808-16>
- Kudhongania, A., & Cordone, A. J. (1974). Batho-spatial distribution pattern and biomass estimate of the major demersal fishes in Lake Victoria. *African Journal of Tropical Hydrobiology and Fisheries*, *3*, 15–31.
- Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, *33*(7): 1870–1874, msw054. <https://doi.org/10.1093/molbev/msw054>
- Louws, F., Rademaker, J., & de Bruijn, F. (1999). The three Ds of PCR-based genomic analysis of phytopathogens: Diversity, detection, and disease diagnosis. *Annual Review of Phytopathology*, *37*, 81–125. <https://doi.org/10.1146/annurev.phyto.37.1.81>
- McAndrew, B. (2000). Evolution, phylogenetic relationships and biogeography. *Fish and Fisheries Series*, *25*, 1–32.
- Molnar, J. L., Gamboa, R. L., Revenga, C., & Spalding, M. D. (2008). Assessing the global threat of invasive species to marine biodiversity. *Frontiers in Ecology and the Environment*, *6*, 485–492. <https://doi.org/10.1890/070064>
- Munang'andu, H. M., Fredriksen, B. N., Mutoloki, S., Brudeseth, B., Kuo, T.-Y., Marjara, I. S., ... Evensen, Ø. (2012). Comparison of vaccine efficacy for different antigen delivery systems for infectious pancreatic necrosis virus vaccines in Atlantic salmon (*Salmo salar* L.) in a cohabitation challenge model. *Vaccine*, *30*, 4007–4016. <https://doi.org/10.1016/j.vaccine.2012.04.039>
- Munang'andu, H. M., Fredriksen, B. N., Mutoloki, S., Dalmo, R. A., & Evensen, Ø. (2013). The kinetics of CD4+ and CD8+ T-cell gene expression correlate with protection in Atlantic salmon (*Salmo salar* L.) vaccinated against infectious pancreatic necrosis. *Vaccine*, *31*, 1956–1963. <https://doi.org/10.1016/j.vaccine.2013.02.008>
- Munang'andu, H. M., Mutoloki, S., & Evensen, O. (2016). Prevention and Control of Viral Diseases in Aquaculture. In F. S. B. Kibenge & M. G. Godoy (Eds.), *Aquaculture virology* (p. 17). London, UK: Academic Press, Elsevier.
- Munang'andu, H. M., Sandtrø, A., Mutoloki, S., Brudeseth, B. E., Santi, N., & Evensen, Ø. (2013). Immunogenicity and cross protective ability of the central VP2 amino acids of infectious pancreatic necrosis virus in Atlantic salmon (*Salmo salar* L.). *PLoS ONE*, *8*, e54263. <https://doi.org/10.1371/journal.pone.0054263>
- Nicholson, P., Fathi, M., Fischer, A., Mohan, C., Schieck, E., Mishra, N., ... Jores, J. (2017). Detection of Tilapia Lake Virus in Egyptian fish farms

- experiencing high mortalities in 2015. *Journal of Fish Diseases*, 40, 1925–1928. <https://doi.org/10.1111/jfd.12650>
- Njiru, M., Nyamweya, C., Gichuki, J., Mugidde, R., Mkumbo, O., & Witte, F. (2012). Increase in anoxia in Lake Victoria and its effects on the fishery. In P. Padilla (Ed.), *Anoxia* (p. 99–128). Rijeka, Croatia: InTech.
- Ogutu-Ohwayo, R. (1990). The decline of the native fishes of lakes Victoria and Kyoga (East Africa) and the impact of introduced species, especially the Nile perch, *Lates niloticus*, and the Nile tilapia, *Oreochromis niloticus*. *Environmental Biology of Fishes*, 27, 81–96. <https://doi.org/10.1007/BF00001938>
- Ogutu-Ohwayo, R. (1994). Adjustments in fish stocks and in life history characteristics of the Nile perch, *Lates niloticus* L. in lakes Victoria, Kyoga and Nabugabo.
- Palese, P., & Schulman, J. L. (1976). Mapping of the influenza virus genome: Identification of the hemagglutinin and the neuraminidase genes. *Proceedings of the National Academy of Sciences*, 73, 2142–2146. <https://doi.org/10.1073/pnas.73.6.2142>
- Surachetpong, W., Janetanakit, T., Nonhabenjawan, N., Tattiyapong, P., Sirikanchana, K., & Amonsin, A. (2017). Outbreaks of Tilapia Lake Virus Infection, Thailand, 2015–2016. *Emerging Infectious Diseases*, 23, 1031. <https://doi.org/10.3201/eid2306.161278>
- Tal, S., Bornestein, B., Ben-Moshe, E., Nevo, Y., Bronstein, M., Bar, N., ... Dishon, A. (2016). Isolation and primary characterization of a virus causing mortality in tilapia. Unpublished NCBI KU552132.1 (Research and Development, KoVax Ltd., 19 Hartom ST., Jerusalem 45212, Israel).
- Tamura, K., & Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, 10, 512–526.
- Tattiyapong, P., Dachavichitlead, W., & Surachetpong, W. (2017). Experimental infection of Tilapia Lake Virus (TiLV) in Nile tilapia (*Oreochromis niloticus*) and red tilapia (*Oreochromis spp.*). *Veterinary Microbiology*, 207, 170–177. <https://doi.org/10.1016/j.vetmic.2017.06.014>
- Tsofack, J. E. K., Zamostiano, R., Watted, S., Berkowitz, A., Rosenbluth, E., Mishra, N., ... Ferguson, H. (2016). Detection of Tilapia Lake Virus (TiLV) in Clinical Samples by Culturing and Nested RT-PCR. *Journal of Clinical Microbiology*, 55, 759–767.
- Witte, F., Goldschmidt, T., Goudswaard, P., Ligtvoet, W., van Oijen, M., & Wanink, J. (1991). Species extinction and concomitant ecological changes in Lake Victoria. *Netherlands Journal of Zoology*, 42, 214–232. <https://doi.org/10.1163/156854291X00298>

How to cite this article: Mugimba KK, Chengula AA, Wamala S, et al. Detection of tilapia lake virus (TiLV) infection by PCR in farmed and wild Nile tilapia (*Oreochromis niloticus*) from Lake Victoria. *J Fish Dis*. 2018;41:1181–1189. <https://doi.org/10.1111/jfd.12790>